



Haringey

Clinical Commissioning Group

Common Mental Health Disorders

This profile has been developed to support the Mental Health Intelligence Network's aim to improve the health of communities by making data and information accessible.

The Common Mental Disorders profile presents all CCG level indicators contained within the Common Mental Health Disorders Profiling Tool: <http://fingertips.phe.org.uk/profile-group/mental-health/profile/common-mental-disorders> The tool provides commissioners, service providers, clinicians, services users and their families with the means to benchmark their area against similar populations and gain intelligence about what works.

Indicators are presented under the headings Risk and Related Factors, Prevalence, Services, Quality and Outcomes, and Finance.

Note: this profile presents CCG geography indicators only and therefore only shows part of the picture (in particular for Risk and Prevalence where many indicators are local authority based). To see all indicators please view the Common Mental Health Disorders Profiling Tool.

Risk and related factors

	Period	Local value	Eng. value	Eng. lowest	Range	Eng. highest
Socioeconomic deprivation: overall IMD score	2012	35.6	21.5	5.8		47.4
Older people living in income deprived households: % of people over 60	2012	36.1	18.1	7.4		56.2
People with CHD and/or diabetes: % patients on the GP register	2012/13	5.7	7.3	4.3		10.4

Prevalence

	Period	Local value	Eng. value	Eng. lowest	Range	Eng. highest
People estimated to have any common mental health disorder: Estimated % of population aged 16-74	2014/15	17.57 ^	15.62	10.29		25.51
Adults with depression known to GPs: Patients with depression as % of all patients on the GP register	2012/13	3.7	5.8	2.9		11.5
New cases of depression: Adults with a new diagnosis of depression as % of all patients on the GP register	2012/13	0.7	1.0	0.5		1.9
Long term mental health problems among GP survey respondents: % people completing GP patient survey who report long-term mental health problem	2013/14	4.6	4.6	2.6		8.8
Depression and anxiety among GP survey respondents: % of people completing GP patient survey reporting they feel moderately or extremely anxious or depressed	2013/14	13.6	12.1	7.2		19.4

Note: ^ - Value estimated

Services

		Period	Local value	Eng. value	Eng. lowest	Range	Eng. highest
Adults with depression known to GPs: Patients with depression as % of all patients on the GP register	■	2012/13	3.7	5.8	2.9		11.5
New cases of depression: Adults with a new diagnosis of depression as % of all patients on the GP register	■	2012/13	0.7	1.0	0.5		1.9
Antidepressant prescribing: Average daily quantities (ADQs) per STAR-PU	■	2013/14	0.7	1.2	0.5		1.8
Use of '1st choice' antidepressants: % of prescription items that were '1st choice' generic SSRIs	■	2013/14	62.6	63.5	53.9		74.6
Hypnotics prescribing: Average daily quantities (ADQs) per STAR-PU	■	2013/14	0.75	1.16	0.53		2.21
IAPT referrals: Rate (quarterly) per 100,000 population aged 18+	■	2013/14 Q4	747	708	6		2136
IAPT referrals for depression: Rate (quarterly) per 100,000 population aged 18+	■	2013/14 Q4	107	82	2		1084
IAPT referrals for mixed anxiety and depression: Rate (quarterly) per 100,000 population aged 18+	■	2013/14 Q4	87.2	83.7	1.2		760.0
IAPT use by BME groups: % of referrals (in quarter) which are for people of black and minority ethnic groups	■	2013/14 Q4	59.9	16.3	0.0		73.5
Access to IAPT services: People entering IAPT (in month) as % of those estimated to have anxiety/depression	■	Dec 2013	8.7	9.5	0.0		50.4
Entering IAPT treatment: Rate (quarterly) beginning IAPT treatment per 100,000 population aged 18+	■	2013/14 Q4	496	469	13		1295
Completion of IAPT treatment: Rate (quarterly) completing treatment per 100,000 population aged 18+	■	2013/14 Q4	289	242	5		611
IAPT diagnosis coding completeness: % (in quarter) of IAPT referrals with a provisional diagnosis code	■	2013/14 Q4	52.7	44.1	0.0		99.3

Quality and Outcomes

		Period	Local value	Eng. value	Eng. lowest	Range	Eng. highest
Assessment of depression: % of adults with a new diagnosis of depression with assessment of severity at treatment outset	■	2012/13	74.2	79.8	62.3		94.4
Follow-up assessment of depression: % of adults with a new diagnosis of depression with follow-up assessment after 4-12 weeks	■	2012/13	55.6	56.0	44.4		71.3
Case finding for depression: % of patients on diabetes and/or CHD register for whom case finding for depression has been undertaken during the preceding 15 months	■	2012/13	84.4	85.9	80.2		91.7
Exception rate for depression: % of patients on depression register excluded from quality indicators	■	2012/13	5.1	5.3	3.1		9.7
Support for people with LTCs: % of people with long term conditions visiting GP who feel they have had enough support from local services in last 6 months	■	2012/13	55.8	64.0	54.3		71.0
Waiting < 28 days for IAPT: % of referrals (in month) waiting	■	Mar 2014	81.7	62.5	4.0		100
Waiting > 90 days for IAPT: % of referrals (in month) waiting > 90 days for first treatment	■	Mar 2014	6.7	10.8	1.0		80.0
IAPT recovery: % of people (in month) who have completed IAPT treatment who are "moving to recovery"	■	Mar 2014	39.4	45.5	24.6		80.0
IAPT reliable recovery: % of people (in quarter) who have completed IAPT treatment who achieved "reliable improvement"	■	2013/14 Q4	64.9	61.9	28.6		75.5
IAPT DNAs: % of IAPT appointments (in quarter) where patient did not attend and gave no advance warning	■	2013/14 Q4	9.7	11.9	1.9		28.6
Patients finishing a course of treatment: % of patients entering IAPT service who receive a course of treatment	■	2013/14 Q4	55.0	51.0	7.2		100
Paired data completeness: % of completed treatments (in quarter) with paired PHQ9 and ADSM scores	■	2013/14 Q4	98.3	96.6	69.7		100

Finance

	Period	Local value	Eng. value	Eng. lowest	Range	Eng. highest
Specialist mental health services spend: rate (£000s) per 100,000 aged 18+ (mapped from PCT)	2012/13	33167	26756	14296		49755
% spend on specialist mental health services: % of all secondary care service spend categorised as mental health (mapped from PCT)	2012/13	14.9	11.9	8.1		19.1
Spend on other mental health services: rate (£000s) per 100,000 population aged 18+ (mapped from PCT)	2012/13	20959	13772	3903		30893
% spend on other mental health: % of all mental health spend categorised as other mental health (mapped from PCT)	2012/13	63.2	51.5	13.1		87.7
Primary care prescribing spend on other mental health: rate (£000s) per 100,000 aged 18+ (mapped from PCT)	2012/13	491	857	268		1419
Secondary Care spend on other mental health: rate (£000s) per 100,000 aged 18+ (mapped from PCT)	2012/13	12425	6762	103		27077
Community care spend on other mental health: rate (£000s) per 100,000 aged 18+ (mapped from PCT)	2012/13	1468	2739	0		15718
Spend on Psychological Therapy Services (IAPT): rate (£000s) per 100,000 aged 16 - 64 (mapped from PCT)	2010/11	326	487	1		1621
Spend on Psychological Therapy Services (Non IAPT): rate (£000s) per 100,000 aged 16 - 64 (mapped from PCT)	2010/11	742	534	11		1809
Cost of GP prescribing for antidepressant drugs: Net Ingredient Cost (£) per 1,000 STAR-PU (quarterly)	2013/14 Q4	26.4	54.4	15.6		163.5
Cost of GP prescribing for hypnotics and anxiolytics: Net Ingredient Cost (£) per 1,000 STAR-PU (quarterly)	2013/14 Q4	78.4	155.0	38.9		497.3


CMHD pathway

	Period	Local value	Eng. value	Eng. lowest	Range	Eng. highest
Socioeconomic deprivation: overall IMD score	2012	35.6	21.5	5.8		47.4
Long-term health problems or disability: % of people whose day-to-day activities are limited by their health or disability	2011	14.0	17.6	11.2		25.6
People estimated to have any common mental health disorder: Estimated % of population aged 16-74	2014/15	17.57 ^	15.62	10.29		25.51
Adults with depression known to GPs: Patients with depression as a % of all patients on the GP register	2012/13	3.7	5.8	2.9		11.5
New cases of depression: Adults with a new diagnosis of depression as % of all patients on the GP register	2012/13	0.7	1.0	0.5		1.9
Assessment of depression: % of adults with a new diagnosis of depression with assessment of severity at treatment outset	2012/13	74.2	79.8	62.3		94.4
Antidepressant prescribing: Average daily quantities (ADQs) per STAR-PU	2013/14	0.7	1.2	0.5		1.8
Access to IAPT services: People entering IAPT services as a % of those estimated to have anxiety/depression	Dec 2013	8.7	9.5	0.0		50.4
Waiting < 28 days for IAPT: % of referrals (in month) waiting	Mar 2014	81.7	62.5	4.0		100
Completion of IAPT treatment: Rate (quarterly) completing treatment per 100,000 population aged 18+	2013/14 Q4	289	242	5		611

Note: ^ - Value estimated

CMHD pathway continued	Period	Local value	Eng. value	Eng. lowest	Range	Eng. highest
IAPT diagnosis coding completeness: % (in quarter) of IAPT referrals with a provisional diagnosis code	2013/14 Q4	52.7	44.1	0.0		99.3
Patients finishing a course of treatment: % of patients entering IAPT service who receive a course of treatment	2013/14 Q4	55.0	51.0	7.2		100
Paired data completeness: % of completed treatments (in quarter) with paired PHQ9 and ADASM scores	2013/14 Q4	98.3	96.6	69.7		100
Rate of recovery: % people "moving to recovery" of those who have completed IAPT treatment	Mar 2014	39.4	45.5	24.6		80.0
IAPT reliable recovery: % of people (in quarter) who have completed IAPT treatment who achieved "reliable improvement"	2013/14 Q4	64.9	61.9	28.6		75.5
Spend on Psychological Therapy Services (IAPT and non IAPT): rate (£000s) per 100,000 aged 16 - 64 (mapped from PCT)	2010/11	1069	1021	15		2649
Secondary care contacts for Common Mental Health Disorders: Rate per 100,000 population aged 18+	2014/15 Q1	466	532	0		2352

- Significantly lower than England average
- Not significantly different from England average
- Significantly higher than England average
- Significance not calculated
- ◆ Regional average



Data quality: ■ Robust ■ Some concerns ■ Significant concerns

Indicators included in spine-charts are drawn from a range of sources, are based on differing populations and are presented for a number of time periods. Detail relating to each indicator is included within the tool under 'definitions'.

<http://fingertips.phe.org.uk/profile-group/mental-health/profile/common-mental-disorders>